



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of a particular form, treating it as a subspecies in one paper and a full species in the next. In the case of the Texas mole described as a subspecies by Dr. J. A. Allen in 1891 and raised to specific rank by the same author in 1893, Dr. Trouessart adopts a curious course. He gives it as a full species with 1893 as the date, and then in synonymy gives the subspecific form in which it was originally described, with 1891 as the date, showing that he was aware of the correct date. Of course, the species should date from 1891—the year in which *the animal* was named—for the date on which an author happens to change his mind as to the rank of a particular form has nothing to do with the date of the name. If this case represents Dr. Trouessart's views in this matter the inference is that he, like some botanists of the old school, is a worshipper of the 'combination.' He certainly agrees with these botanists in spelling personal and some other specific names with a capital initial letter—in this respect again departing from the best usage among zoologists.

Sections of genera and forms of species of earlier authors are sometimes given formal subgeneric and subspecific names, and names so given are credited to the early author instead of to himself. Thus the section 'Cercopithecini Rhinosticti' of Sclater is made the subgenus '*Rhinostictus* Sclater,' and Dr. Harrison Allen's 'Var. (b) Northern form of *Vespertilio gryphus*' is made 'Var. *septentrionalis* H. Allen.'

I am indebted to Dr. T. S. Palmer for calling my attention to Dr. Trouessart's extraordinary rule for the treatment of preoccupied names. If he finds such names preoccupied among mammals he promptly renames them (as *Scaptogale* nob. for *Echinogale* Pomel), but if they are preoccupied in other branches of the animal kingdom he lets them stand. Thus the generic names *Tylostoma* (p. 155), *Schizostoma* (154), *Macrotus* (152), *Mystacina* (149), *Furia* (135), *Vesperus* (106), *Megaloglossus* (98) and many others are retained, notwithstanding the fact that all are preoccupied and replaced by other names now in more or less common use. A few of the dates given for genera are erroneous. For instance, *Leuconoe* Boie '1825' should be 1830, and *Dendrogale* Gray '1843' should be

1848. *Prototalpa* is evidently an amended form of *Protalpa* and as such should date from *Prototalpa* Roger 1887 instead of *Protalpa* Filhol 1877.

Since the appearance of the first edition of Dr. Trouessart's Catalogue (1878-85) no attempt has been made to collect in one work the names of all the mammals of the world; and since all fossil as well as living species are included, the immensity of the task is apparent. Most authors shrink from such an undertaking, not only on account of its magnitude, but also on account of the extreme difficulty, not to say impossibility, of determining the status of described forms in groups that have not been recently revised. Nevertheless the work is of such great practical utility that for years to come every student of living or fossil mammals must keep a copy at his elbow and will owe its author a debt of gratitude. Dr. Trouessart is evidently a very rapid worker; we wish him health and freedom from interruptions, so that his great undertaking may be speedily completed.

C. H. M.

SOCIETIES AND ACADEMIES.

THE 96TH REGULAR MEETING OF THE CHEMICAL SOCIETY OF WASHINGTON.

THE following program was presented:

H. N. Stokes: 'The Chloronitrides of Phosphorus.' P. Fireman: 'The Ripening of Cheese and the Rôle which Micro-organisms Play in the Process.' E. A. de Schweinitz and Marion Dorset: 'The Product of the Tuberculosis Bacillus.' H. W. Wiley and W. H. Krug: 'The Standard Methods of Starch Determination.' W. H. Krug and J. E. Blomén: 'The Commercial Preparation of Nitro-naphthalene.' F. K. Cameron: 'The Replacement of Chlorine by Sulphur in Alkaline Chlorides.' Wirt Tassin: 'A New Mineral.'

Dr. Stokes showed that the only members of the phosphorus Chloronitrides series $(\text{PNCl}_2)_n$, hitherto known are $(\text{PNCl}_2)_3$ (Liebig) and $(\text{PNCl}_2)_4$ (Stokes). The series is now extended to include $(\text{PNCl}_2)_5$, $(\text{PNCl}_2)_6$ and $(\text{PNCl}_2)_7$, as well as a mixture of higher polymers, not yet isolated, and terminating with a rubber-like polymer of high molecular weight. The substances are prepared by heating equimolecular weights of phosphorus pentachloride

and ammonium chloride in sealed tubes, whereby a mixture of chlorides in nearly theoretical amount is obtained. This is distilled from the open tube and afterwards submitted to careful fractional distillation *in vacuo*. The unique feature of the series is found in the fact that any member can be converted into the rubber-like polymer by heating, and that the latter, on distilling at a higher temperature, breaks down quantitatively into a mixture of all the lower members; it is, therefore, possible ultimately to convert any member completely into any other by heating and distilling alone. The lower members, up to and including $(\text{PNCl}_2)_6$ are well characterized, finely crystallized bodies, while $(\text{PNCl}_2)_7$ is liquid. Their stability diminishes with increasing molecular weight, $(\text{PNCl}_2)_8$ being unattached by boiling alkalis, while the rubber-like polymer is destroyed by boiling water. The formation of PNCl_2 and $(\text{PNCl}_2)_2$ could not be detected. The chloronitrides constitute the first extended series of inorganic polymers interconvertible by simple and direct means.

Dr. Fireman gave an extended résumé of the work of Duclaux and others showing that the lactic acid producing bacteria are not so important in the production of butter flavors as are the peptonizing bacteria.

Drs. de Schweinitz and Dorset stated that in studying the products of the growth of the tuberculosis bacilli in artificial media, it was noted that the reaction of the cultures usually becomes acid, and as Prudden and Hodenpyl had succeeded in producing tuberculous nodules without necrosis by the intravenous injection of dead bacilli, it seemed as though it should be possible to isolate from cultures of tuberculosis bacilli an acid substance, which is responsible for the necrosis of tissue that always takes place in this disease. After many fruitless attempts they succeeded in isolating from artificial cultures a crystalline acid substance having a melting point of 161° to 164° C. which was soluble in ether, alcohol and water, and crystallized in needle-like prisms. The solution of this substance was optically inactive and did not give the biuret action. The preliminary analysis of this substance gave a formula, like that of teraconic acid and the other

properties correspond closely with this acid. Its identity if such has not yet been proved. When injected subcutaneously into guinea pigs it causes a slight inflammation and localized necrosis, and injected directly into the liver tissue by means of a hypodermic syringe it produces characteristic necrosis. The substance causes a reduction of temperature in tuberculous animals, and it seems probable that we have here the material which is responsible for the necrosis in tubercular infection.

Dr. Wiley and Mr. Krug showed that various methods depending on the polarization of the starch or its inversion products, as well as others based on direct weighing of the starch, were too inaccurate to commend themselves to the analyst. The only accurate methods depend on the eventual inversion of the starch into dextrose, which is then estimated by Allihn's method. Experiments in the preparation of the Nitro-naphthalenes showed that they can not be prepared from α -naphthalene sulphonic acid, but that the best yield is obtained with mixtures of nitric and sulphuric acids. The best results were obtained with 30° B. nitric acid, using about three times as much acid as naphthalene. The amount of sulphuric acid, used varies with the nitration degree desired, varying from 4:1 (nitric: sulphuric) for low melting products to 3:2 for the higher derivations.

Dr. Cameron cited some of his own experiments to show that, although it is possible in the case of some heavy metals to replace chlorine by sulphur by merely boiling the latter with a solution of the chloride, it is impossible to obtain such a reaction with alkaline, or alkaline earth chlorides even upon heating in sealed tubes at a high temperature.

Mr. Tassin gave a short description of a new mineral, but reserves the details for special publication.

V. K. CHESNUT.

Secretary.

TORREY BOTANICAL CLUB, MAY 26, 1897.

THE President, Hon. Addison Brown, presided. The evening was devoted to a lecture by Mrs. Elizabeth G. Britton, entitled 'The

Moss Flora of the Adirondack Mountains,' illustrated by lantern-slides prepared by Mr. C. H. Van Brunt, and also by numerous mounted specimens. These specimens, comprising about 150 sheets, handsomely displayed about the walls of the lecture-room, represented collections made by Mrs. Britton in the vicinity of Adirondack Lodge and Lake Placid in the years 1892, 1894 and 1896. The various locations where these mosses grew were described, including the story of a climb up Mount Whiteface. Graphic bits of description of these mosses brought out salient points as the slides indicating their structure were exhibited, and these were accompanied by numerous slides illustrating the scenery of their habitat. Among about 30 rare species enumerated were *Raphidostegium Jamesii*, not previously reported for New York State, and *Bryum concinatum*, found only once before in the United States. Duplicates of Mrs. Britton's collection have been deposited at the State Herbarium at Albany, the main collection having been presented to the Herbarium of Columbia University. Partial sets were sent to the Brooklyn Institute, to Cornell University and to various other collections.

The subject of the lecture was further discussed by Mr. A. P. Grout and by Mrs. Britton, after which the Club adjourned to the second Tuesday in October, field meetings continuing meantime on Saturdays.

EDWARD S. BURGESS,
Secretary.

SCIENCE CLUB OF THE UNIVERSITY OF WISCONSIN.

MR. EDWARD KREMER discussed 'The Periodic System.' The speaker introduced his subject by giving a brief account of the earlier attempts to refer back to a prime matter the manifold forms in which material nature reveals herself to man. This line of thought was briefly traced from the early Greek philosophers to Prout. The reaction that followed Prout's hypothesis was duly emphasized. The lack of interest in Doehereiner's trials, and the modifications of these trials by others, in Newland's octaves, in Lothar Meyer's first chart it was thus explained in large part. Attention was also called to the fact that the principal interests

seemed to center about the numerical relation existing between the atomic weights of the elements and that the time value of the system was not duly appreciated until this was made secondary to the natural classification of the elements according to their physical and chemical properties. Incidentally the prediction by Mardelejeff of the existence of certain elements and their discovery by others were alluded to, because with the discovery of their elements the natural law gained more general favor. What the periodic law really is was strongly emphasized, also the fact that the present arrangement of the elements according to this law is imperfect, but more recent systems were not discussed, partly for want of time. Finally the great advantage to be gained by using the periodic law as the basis for teaching advanced chemistry was demonstrated.

C. S. SLICHTER reviewed Larmor's 'Theory of the Ether.' He first gave an explanation of the vortex theory of matter, including the properties of vortex rings and their mode of linking, and kinds of vibration. Hill's spherical vortex was explained by a diagram, and the possibility of building up a rotationally elastic ether from such cells was explained. Photographs of sandstone showing turbulent motion were presented for the purpose of showing the character of their structure. The intrinsic energy of the ether, radiation, action at a distance, density and inertia of common matter were then discussed, closing with a new theory of gravitation. Mr. Quantz gave the results of work on the psychology of rapid reading. They were the results of a statistical study of the mental habits and characteristics of fifty university students, showing that rapidity of reading depends chiefly upon the following factors, given in the order of their importance: (a) quickness of visual perception; (b) practice, as measured by extent of reading from childhood up; (c) power of mental concentration; (d) absence of motor-mindedness (lip movement in silent reading); (e) rapidity of mental operations in general; (f) strongly developed eye-mindedness; (g) scholarly ability, as determined by class study.

WM. S. MARSHALL,
Secretary.

THE ACADEMY OF SCIENCE OF ST. LOUIS.

At the meeting of the Academy of Science of St. Louis on the 7th of June, 1897, twenty-one persons present, Mr. Robert Combs, of Ames, Iowa, presented a paper entitled 'Plants Collected in the District of Cienfuegos, Province of Santa Clara, Cuba, in 1895-1896.' The paper embraces the results of a collection extending from the commencement of the rainy season of one year until the close of the dry season the following spring, the territory covered by the collection lying between the entrance of the Bay of Cienfuegos, on the south coast of Cuba, up the bay and the river Damuji to Rodas, and extending back from the river to Yaguaramos and almost to the Cienega de Zapato, a region including nearly all kinds of soil and conditions found upon the island, except those of the mountain regions and the mud swamps. A brief statement was made concerning the origin of the Cuban flora and its affinities with that of continental Central America, rather than the geographically nearer Floridan region.

The paper comprised a full catalogue of the collections made, which had been determined at the herbarium of Harvard University, and of which several sets had been distributed to the larger herbaria.

Professor F. E. Nipher made some remarks on the difficulties yet involved in the theories of the ether.

WILLIAM TRELEASE.

THE TEXAS ACADEMY OF SCIENCE.

At the annual meeting of the Texas Academy of Science, held in the chemical lecture room of the University of Texas, June 15th, the following papers were presented:

'The Personality of a Great Genius—Sylvester,' by Dr. George Bruce Halstead, in which the author took occasion to point out the exalted position of the higher pure mathematics and to give his estimate of Sylvester—his old friend and teacher—as a mathematician.

'Epsom Salts, Magnesium Sulphate, from Brown county, Texas,' by Dr. H. W. Harper. The occurrence of Epsomite in large quantities in Brown county, and of a purity sufficient to make it the source of an exceedingly cheap commercial product, was here announced for

the first time. Dr. Harper's analyses gave the following results:

Water.....	40.07	40.00
Silica	21.075	21.43
Alumina and Iron Oxides..	2.20	2.21
Magnesium Oxide	12.381	12.38
Calcium Oxide.....	trace.	
Sulphur Trioxide.....	24.014	24.01
	99.74	100.03

Calculated to contain 76.13 % $\text{MgSO}_4 \cdot 7 \text{H}_2\text{O}$.

A series of experiments demonstrated that the crude material yields crystallized MgSO_4 within one or two % of the analytical results.

'Some Texas Oil Horizons,' by Hon. E. T. Dumble. The Oil Horizons in Texas range from the Carboniferous to recent formations. I. The lowest horizon in the State is in the Bend division of the Carboniferous—the black shales of the Colorado river near Brownwood. They are apparently a continuation of the asphalt deposits of Indian Territory, as indicated by the finding of similar fossils. II. The lower beds of the Cretaceous—the Trinity Sands—in Jack, Montague and Burnett counties. III. The Eagle Ford Shale (Benton) of the Cretaceous, as, for example, in the well at Fiskeville, which furnished a little oil, and in the railroad cut south of Austin, where, it is said, the shale will burn. IV. The Black Marls of the Black Waxy Prairie Division of the Cretaceous. The well at Corsicana encounters at 1,040 feet a 15-foot oil sand. Oil will also be found eastward to Smith and Anderson counties. V. The Nacogdoches horizon, which seems to be the eastern appearance of the oil and asphalt deposits of California.

'Pedagogical Note on Mensuration,' by Professor Arthur Lefevre.

'The Texas Permian,' by W. F. Cummins (read by title).

'On the Bio-Geography of Mexico and the Southwestern United States,' by C. H. Tyler Townsend (read by title).

The following officers were elected for the ensuing year: President, Dr. George Bruce Halsted; Vice-President, Professor T. U. Taylor; Treasurer, Hon. E. T. Dumble; Secretary, Professor W. W. Norman.

FREDERIC W. SIMONDS.